



BAVENCIO[®]
avelumab Injection
20 mg/mL

PATIENT WALLET CARD

Carry this card with you while on treatment with **BAVENCIO (avelumab)**. Be sure to tell all healthcare providers you see that you are being treated with BAVENCIO and **show them this card.**

Manufactured by: EMD Serono, Inc.
One Technology Place, Rockland, MA
02370 USA.

Marketed by: EMD Serono, Inc.
and Pfizer Inc, NY, NY 10017 USA.

BAVENCIO is a trademark of Merck KGaA,
Darmstadt, Germany.

For more information,
call toll-free 1-844-826-8371 or go to
www.bavencio.com.

**EMD
SERONO**



Copyright © 2019
EMD Serono, Inc.
All rights reserved.

July 2019
US/AVE/0317/0118(1)

INDICATIONS

BAVENCIO is indicated for the treatment of:

- Adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (MCC)
- Patients with locally advanced or metastatic urothelial carcinoma (UC) who have disease progression during or following platinum-containing chemotherapy, or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy

These indications are approved under accelerated approval based on tumor response rate and duration of response. Continued approval for these indications may be contingent upon verification and description of clinical benefit in confirmatory trials.



What is the most important information I should know about BAVENCIO?

BAVENCIO is a medicine that may treat certain cancers by working with your immune system. BAVENCIO can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life-threatening and can lead to death.

Call or see your healthcare provider right away if you get any symptoms of the following problems or these symptoms get worse:

Lung problems (pneumonitis).

Signs and symptoms of pneumonitis may include:

- new or worsening cough
- shortness of breath
- chest pain

Liver problems, including hepatitis.

Signs and symptoms of liver problems may include:

- yellowing of your skin or the whites of your eyes

- severe nausea or vomiting
- pain on the right side of your stomach-area (abdomen)
- drowsiness
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Intestinal problems (colitis).

Signs and symptoms of colitis may include:

- diarrhea (loose stools) or more bowel movements than usual
- blood in your stools or dark, tarry, sticky stools
- severe stomach-area (abdomen) pain or tenderness

Hormone gland problems (especially the adrenal glands, thyroid, and pancreas). Signs and symptoms that your hormone glands are not working properly may include:

- rapid heart beat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- hair loss

- changes in mood or behavior, such as irritability or forgetfulness
- constipation
- your voice gets deeper
- very low blood pressure
- urinating more often than usual
- dizziness or fainting
- nausea or vomiting
- stomach area (abdomen) pain
- feeling cold

Kidney problems, including nephritis.

Signs and symptoms of kidney problems may include:

- decrease in your amount of urine
- blood in your urine
- swelling in your ankles
- loss of appetite

Problems in other organs. Signs and symptoms may include:

- severe muscle weakness
- severe or persistent muscle or joint pains
- chest pain and tightness
- trouble breathing
- skin rash, blisters, or peeling

- changes in heartbeat, such as beating fast, or seeming to skip a beat, or pounding sensation
- tiredness, sleepiness
- swelling of the feet and legs
- dizziness or fainting
- fever, flu like symptoms
- changes in eyesight

Severe infusion reactions. Signs and symptoms of severe infusion reactions may include:

- chills or shaking
- hives
- flushing
- shortness of breath or wheezing
- low blood pressure
- fever
- back pain
- abdominal pain

Heart problems. When BAVENCIO is used with the medicine axitinib, severe heart problems can happen and can lead to death. Signs and symptoms of heart problems may include:

- swelling of your stomach-area, legs, hands, feet or ankles
- shortness of breath
- nausea or vomiting
- chest discomfort, including pain or pressure
- weight gain
- pain or discomfort in your arms, back, neck, or jaw
- breaking out in a cold sweat
- feeling lightheaded or dizzy

Getting medical treatment right away may help to keep these problems from becoming more serious.

Your healthcare provider will check you for these problems during your treatment with BAVENCIO. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may delay or completely stop treatment with BAVENCIO if you have severe side effects.

What is BAVENCIO?

BAVENCIO is a prescription medicine used to treat:

- a type of skin cancer called Merkel cell carcinoma (MCC) in adults and children 12 years of age and older. BAVENCIO may be used when your skin cancer has spread.
- a type of cancer in the bladder or urinary tract called urothelial carcinoma (UC). BAVENCIO may be used when your cancer:
 - has spread or cannot be removed by surgery (advanced UC), **and**
 - you have received chemotherapy that contains platinum, and it did not work or is no longer working.

It is not known if BAVENCIO is safe and effective in children under the age of 12. (continues on the reverse)

Please see accompanying full Prescribing Information and Medication Guide. For more information, visit BAVENCIO.com.

Keep your HCP's contact information within easy reach. Simply fill out and fold to fit into your wallet.

For more information, visit BAVENCIO.com.

My oncologist's contact information

Be sure to tell all healthcare providers you see that you are being treated with **BAVENCIO (avelumab)** and show them this card.

My oncologist's contact information:

My name and telephone number:

Emergency contact information:

Other medicines that I am taking:

Other healthcare provider contact information:

Before you receive BAVENCIO, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease or ulcerative colitis
- have had an organ transplant
- have lung or breathing problems
- have liver or kidney problems
- have diabetes
- have heart problems or high blood pressure
- have a high cholesterol level in your blood
- are pregnant or plan to become pregnant. BAVENCIO can harm your unborn baby. If you are able to become pregnant, you should use an effective method of birth control during your treatment and for at least 1 month after the last dose of BAVENCIO.
- are breastfeeding or plan to breastfeed. It is not known if BAVENCIO passes into your breast milk. Do not breastfeed during treatment and for at least 1 month after the final dose of BAVENCIO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive BAVENCIO?

- Your healthcare provider will give you BAVENCIO into your vein through an intravenous (IV) line over 60 minutes.
- BAVENCIO is usually given every 2 weeks.
- Your healthcare provider will give you medicines before the first 4 infusions and then as needed to help reduce infusion reactions.
- Your healthcare provider will decide how many treatments you need.
- Your healthcare provider will do blood tests to check you for certain side effects.
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment.

What are the possible side effects of BAVENCIO?

BAVENCIO can cause serious side effects, including:

- See "What is the most important information I should know about BAVENCIO?"

The most common side effects of BAVENCIO in people with MCC include:

- feeling tired
- muscle and bone pain
- diarrhea
- nausea
- infusion-related reaction including chills, fever and back pain
- rash
- decreased appetite
- swelling in your hands, feet, or ankles

The most common side effects of BAVENCIO in people with UC include:

- feeling tired
- infusion-related reaction including chills, fever, back pain, redness and shortness of breath
- muscle and bone pain
- nausea
- decreased appetite
- urinary tract infection

These are not all the possible side effects of BAVENCIO.

Ask your healthcare provider or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



General information about the safe and effective use of BAVENCIO.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. If you would like more information about BAVENCIO, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about BAVENCIO that is written for health professionals.

What are the ingredients in BAVENCIO?

Active ingredient: avelumab
Inactive ingredients: D-mannitol, glacial acetic acid, polysorbate 20, sodium hydroxide, and Water for Injection

Please see accompanying full Prescribing Information and Medication Guide. For more information, visit BAVENCIO.com.