

Fighting advanced kidney cancer **TOGETHER**

BAVENCIO® (avelumab) with INLYTA® (axitinib)

INDICATION

BAVENCIO is a prescription medicine used to treat a type of kidney cancer called renal cell carcinoma (RCC). BAVENCIO may be used with the medicine INLYTA as your first treatment when your kidney cancer has spread or cannot be removed by surgery (advanced RCC).

It is not known if BAVENCIO and INLYTA are safe and effective in children.



Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for **BAVENCIO** and full [Prescribing Information/Patient Information](#) for **INLYTA**.

 **BAVENCIO**[®]
avelumab injection
20 mg/mL



 **Inlyta**[®]
axitinib imaging tablets



About Advanced Renal Cell Carcinoma (RCC)

When a person has cancer, the body's cells start to grow at abnormal rates with irregular shapes. Sometimes the cells form a mass called a tumor.

As cancer cells grow, some may spread to nearby organs or to lymph nodes, forming what is known as a locally advanced tumor. Cancer cells can also enter the bloodstream and spread to more distant organs, which is called metastasis.

When a tumor starts from kidney cells, it's called kidney cancer. RCC is the most common type of kidney cancer. Nine in 10 people who have kidney cancer have RCC.

Advanced RCC is when kidney cancer has spread or cannot be removed by surgery.



It's important to talk to your healthcare provider about your options and work together to make a treatment plan.

How BAVENCIO® (avelumab) and INLYTA® (axitinib) Work

BAVENCIO and INLYTA are 2 different types of medicines that work in 2 different ways.

BAVENCIO

- BAVENCIO is an immunotherapy—a medicine that may treat advanced RCC by working with your immune system. BAVENCIO can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life-threatening and can lead to death
- BAVENCIO is a man-made antibody. Your body naturally produces antibodies to help your immune system work
- BAVENCIO may engage both your adaptive and innate immune systems
- The immune system is a part of your body that fights diseases to keep you healthy. One job of your immune system is to remove cells that are becoming cancerous. Over time, cancer cells may be able to stop your immune system from attacking them
- Your adaptive immune system learns to identify and adapt to threats over time, and delivers a response to defend against specific threats
- Your innate immune system is your first line of defense against general threats

INLYTA

- In order for cancer cells to spread, they need a constant supply of oxygen and nutrients. They get these from blood vessels
- When a tumor reaches a certain size, it needs new blood vessels to keep growing
- Animal and laboratory studies suggest that INLYTA prevents new blood vessels from forming
- It is believed that this may help stop tumors from growing and cancer from spreading


Indication

BAVENCIO is a prescription medicine used to treat a type of kidney cancer called renal cell carcinoma (RCC). BAVENCIO may be used with the medicine INLYTA as your first treatment when your kidney cancer has spread or cannot be removed by surgery (advanced RCC).

It is not known if BAVENCIO and INLYTA are safe and effective in children.

Serious Side Effects

- **BAVENCIO can cause serious side effects, including:** lung problems (pneumonitis); liver problems, including hepatitis; intestinal problems (colitis); hormone gland problems (especially the adrenal glands, thyroid, and pancreas); kidney problems, including nephritis; problems in other organs; severe infusion reactions; and heart problems.
- **INLYTA may cause serious side effects, including:** high blood pressure (hypertension, which is common and may sometimes be severe); problem with blood clots in your veins or arteries; bleeding; heart failure; tear in your stomach or intestinal wall (perforation); thyroid gland problems; Reversible Posterior Leukoencephalopathy Syndrome (RPLS); protein in your urine; and change in liver function.



Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for BAVENCIO and full [Prescribing Information/Patient Information](#) for INLYTA.



About the BAVENCIO® (avelumab) with INLYTA® (axitinib) Clinical Trial

In a clinical trial of 442 patients taking BAVENCIO with INLYTA versus 444 patients taking a medicine called sunitinib, researchers studied:

- **Progression-free survival (PFS)**, the length of time during and after cancer treatment that a patient lives without disease progression—meaning that the tumors did not grow or spread during this time. This is one way to check how effective a treatment is
- **Overall survival (OS)**, the total time patients on each medicine remained alive after starting treatment
- **Objective response rate (ORR)**, the percentage of patients who no longer had any signs of cancer or whose tumor shrank 30% or more

BAVENCIO with INLYTA is not a cure. The data represent an average of patients in the clinical trial, and not all patients may experience the same results.

Selected Safety Information

The most common side effects of BAVENCIO when given with INLYTA in people with RCC include:

- | | | |
|------------------------|--|-------------------------------|
| • diarrhea | • liver problems | • rash |
| • feeling tired | • blisters or rash on the palms of your hands and soles of your feet | • shortness of breath |
| • high blood pressure | • hoarseness | • cough |
| • muscle and bone pain | • decreased appetite | • stomach-area (abdomen) pain |
| • nausea | • low levels of thyroid hormone | • headache |
| • mouth sores | | |

Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for BAVENCIO and full [Prescribing Information/Patient Information](#) for INLYTA.

Together Can Be Powerful

Study results

In a clinical trial of 442 patients taking BAVENCIO with INLYTA versus 444 patients taking a medicine called sunitinib:




BAVENCIO with INLYTA decreased the overall risk of disease progression or death by 31% compared to sunitinib (progression-free survival).


Patients taking BAVENCIO with INLYTA experienced a median PFS of 13.8 months. Median time period means that half the patients went longer than 13.8 months with no tumor growth or spread, and the other half went less than 13.8 months with no tumor growth or spread. Patients taking sunitinib experienced a median PFS of 8.4 months. That's a 5-month increase in median PFS for patients taking BAVENCIO with INLYTA compared to patients taking sunitinib.

At the time data were collected, overall survival data were immature, which means that there was not yet enough information available to assess the impact of BAVENCIO with INLYTA compared to sunitinib on overall survival.

BAVENCIO with INLYTA reduced tumor size in twice as many patients compared to sunitinib (objective response rate).



BAVENCIO with INLYTA



Sunitinib

- 3.4% of patients (15 of 442) taking BAVENCIO with INLYTA saw their tumors disappear compared to 1.8% of patients (8 of 444) taking sunitinib (complete response)
- 48% of patients (212 of 442) taking BAVENCIO with INLYTA saw their tumors shrink compared to 24% of patients (106 of 444) taking sunitinib (partial response)

This includes patients whose tumors shrank 30% or more and whose response to treatment lasted at least 4 weeks.



Important Safety Information

BAVENCIO® (avelumab)

BAVENCIO is a medicine that may treat certain cancers by working with your immune system. BAVENCIO can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life-threatening and can lead to death.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with BAVENCIO. Your healthcare provider may treat you with corticosteroids or hormone replacement medicines. Your healthcare provider may delay or completely stop treatment with BAVENCIO if you have severe side effects.

Call or see your healthcare provider right away if you get any symptoms of the following problems, or if these symptoms get worse.

BAVENCIO can cause serious side effects, including:

- **Lung problems (pneumonitis)**—signs and symptoms of pneumonitis may include:
 - new or worsening cough
 - shortness of breath
 - chest pain

- **Liver problems, including hepatitis**—signs and symptoms of liver problems may include:
 - yellowing of your skin or the whites of your eyes
 - severe nausea or vomiting
 - pain on the right side of your stomach-area (abdomen)
 - drowsiness
 - dark urine (tea colored)
 - bleeding or bruising more easily than normal
- **Intestinal problems (colitis)**—signs and symptoms of colitis may include:
 - diarrhea (loose stools) or more bowel movements than usual
 - blood in your stools or dark, tarry, sticky stools
 - severe stomach-area (abdomen) pain or tenderness
- **Hormone gland problems (especially the adrenal glands, thyroid, and pancreas)**—signs and symptoms that your hormone glands are not working properly may include:

– rapid heart beat	– constipation
– increased sweating	– your voice gets deeper
– extreme tiredness	– very low blood pressure
– weight gain or weight loss	– urinating more often than usual
– feeling more hungry or thirsty than usual	– dizziness or fainting
– hair loss	– nausea or vomiting
– changes in mood or behavior, such as irritability or forgetfulness	– stomach area (abdomen) pain
	– feeling cold

- **Kidney problems, including nephritis**—signs and symptoms of kidney problems may include:
 - decrease in your amount of urine
 - blood in your urine
 - swelling in your ankles
 - loss of appetite
- **Problems in other organs**—signs and symptoms may include:
 - severe muscle weakness
 - severe or persistent muscle or joint pains
 - chest pain and tightness
 - trouble breathing
 - skin rash, blisters or peeling
 - changes in heartbeat, such as beating fast, or seeming to skip a beat, or pounding sensation
 - tiredness, sleepiness
 - swelling of the feet and legs
 - dizziness or fainting
 - fever, flu like symptoms
 - changes in eyesight

- **Severe infusion reactions**—signs and symptoms of severe infusion reactions may include:
 - chills or shaking
 - hives
 - flushing
 - shortness of breath or wheezing
 - low blood pressure
 - fever
 - back pain
 - abdominal pain
- **Heart problems**—when BAVENCIO is used with the medicine INLYTA, severe heart problems can happen and can lead to death. Signs and symptoms of heart problems may include:
 - swelling of your stomach-area, legs, hands, feet or ankles
 - shortness of breath
 - nausea or vomiting
 - chest discomfort, including pain or pressure
 - weight gain
 - pain or discomfort in your arms, back, neck, or jaw
 - breaking out in a cold sweat
 - feeling lightheaded or dizzy

continued on next page

Important Safety Information (cont'd)

Before you receive BAVENCIO, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease or ulcerative colitis
- have had an organ transplant
- have lung or breathing problems
- have liver or kidney problems
- have diabetes
- have heart problems or high blood pressure
- have a high cholesterol level in your blood
- are pregnant or plan to become pregnant. BAVENCIO can harm your unborn baby. If you are able to become pregnant, you should use an effective method of birth control during your treatment and for at least 1 month after the last dose of BAVENCIO
- are breastfeeding or plan to breastfeed. It is not known if BAVENCIO passes into your breast milk. Do not breastfeed during treatment and for at least 1 month after the final dose of BAVENCIO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

INLYTA® (axitinib)

INLYTA may cause serious side effects, including:

- **High blood pressure (hypertension). High blood pressure is common with INLYTA, and may sometimes be severe.**

Your healthcare provider should check your blood pressure regularly during treatment with INLYTA. If you develop blood pressure problems, your healthcare provider may prescribe medicine to treat your high blood pressure, lower your dose, or stop your treatment with INLYTA

- **Problem with blood clots in your veins or arteries.** INLYTA can cause blood clots which can be serious, and sometimes lead to death. Get emergency help and call your healthcare provider if you get any of the following symptoms:
 - chest pain or pressure
 - pain in your arms, back, neck, or jaw
 - shortness of breath
 - numbness or weakness on one side of your body
 - trouble talking
 - headache
 - vision changes

- **Bleeding.** INLYTA can cause bleeding which can be serious, and sometimes lead to death. Call your healthcare provider right away or get medical help if you develop any of the following signs or symptoms:
 - unexpected bleeding or bleeding that lasts a long time, such as:
 - unusual bleeding from the gums
 - menstrual bleeding or vaginal bleeding that is heavier than normal
 - bleeding that is severe or you cannot control
 - pink or brown urine
 - red or black stools (looks like tar)
 - bruises that happen without a known cause or get larger
 - cough up blood or blood clots
 - vomit blood or your vomit looks like “coffee grounds”
 - unexpected pain, swelling, or joint pain
 - headaches, feeling dizzy or weak
- **Heart failure.** Your healthcare provider should check you for signs or symptoms of heart failure regularly during treatment with INLYTA. Heart failure can be serious and can sometimes lead to death. Tell your healthcare provider if you have any of the following symptoms during your treatment with INLYTA:
 - tiredness
 - swelling of your stomach-area (abdomen), legs or ankles
 - shortness of breath
 - protruding neck veins

- **Tear in your stomach or intestinal wall (perforation).** A tear in your stomach or intestinal wall can be serious and can sometimes lead to death. Get medical help right away if you get the following symptoms:
 - severe stomach-area (abdominal) pain or stomach-area pain that does not go away
 - vomit blood
 - red or black stools
- **Thyroid gland problems.** Your healthcare provider should do blood tests to check your thyroid gland function before and during your treatment with INLYTA. Tell your healthcare provider if you have any of the following symptoms during your treatment with INLYTA:
 - tiredness that worsens or does not go away
 - feeling hot or cold
 - your voice deepens
 - weight gain or weight loss
 - hair loss
 - muscle cramps and aches

continued on next page

Important Safety Information (cont'd)

- **Risk of wound healing problems.** Wounds may not heal properly during INLYTA treatment. Tell your healthcare provider if you plan to have any surgery before starting or during treatment with INLYTA.
 - You should stop taking INLYTA at least 2 days before planned surgery
 - Your healthcare provider should tell you when you may start taking INLYTA again after surgery
- **Reversible Posterior Leukoencephalopathy Syndrome (RPLS).** A condition called reversible posterior leukoencephalopathy syndrome (RPLS) can happen while taking INLYTA. Call your healthcare provider right away if you get:
 - headache
 - seizures
 - weakness
 - confusion
 - high blood pressure
 - blindness or change in vision
 - problems thinking
- **Protein in your urine.** Your healthcare provider should check your urine for protein before and during your treatment with INLYTA. If you develop protein in your urine, your healthcare provider may decrease your dose of INLYTA or stop your treatment
- **Change in liver function.** Your healthcare provider should do blood tests before and during your treatment with INLYTA to check your liver function

Before taking INLYTA, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure
- have thyroid problems
- have liver problems
- have a history of blood clots in your veins or arteries (types of blood vessels), including stroke, heart attack, or change in vision
- have any bleeding problems
- have a history of heart failure
- have an unhealed wound
- plan to have surgery or have had a recent surgery. You should stop taking INLYTA for at least 2 days before planned surgery
- have any other medical conditions

For females, tell your healthcare provider if you:

- are pregnant or plan to become pregnant. Taking INLYTA during pregnancy can harm your unborn baby. You should not become pregnant while taking INLYTA. Talk to your healthcare provider if you are pregnant or plan to become pregnant
- are able to become pregnant. You should have a pregnancy test before you start treatment with INLYTA. Use effective birth control during treatment and for at least 1 week after your last dose of INLYTA. Talk to your healthcare provider about birth control methods that you can use to prevent pregnancy during this time
- are breastfeeding or plan to breastfeed. It is not known if INLYTA passes into your breast milk. Do not breastfeed during treatment and for at least 2 weeks after your last dose of INLYTA

For males with female partners who are able to become pregnant:

- use effective birth control during treatment and for at least 1 week after your last dose of INLYTA
- if your female partner becomes pregnant during your treatment with INLYTA, tell your healthcare provider right away

INLYTA may cause **fertility problems** in males and females, which may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. INLYTA and certain other medicines can affect each other causing serious side effects.

Talk with your healthcare provider before you start taking any new medicine. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

Do not drink grapefruit juice or eat grapefruit. Grapefruit may increase the amount of INLYTA in your blood.

The most common side effects of BAVENCIO® (avelumab) when given with INLYTA® (axitinib) in people with RCC include:

- | | |
|--|---------------------------------|
| • diarrhea | • hoarseness |
| • feeling tired | • decreased appetite |
| • high blood pressure | • low levels of thyroid hormone |
| • muscle and bone pain | • rash |
| • nausea | • shortness of breath |
| • mouth sores | • cough |
| • liver problems | • stomach-area (abdomen) pain |
| • blisters or rash on the palms of your hands and soles of your feet | • headache |

These are not all the possible side effects of BAVENCIO in combination with INLYTA. Ask your healthcare provider or pharmacist for more information. You may report side effects to the FDA at **1-800-FDA-1088**, or **www.fda.gov/medwatch**.

Tips to Help Manage Certain Side Effects

The tips on the following pages are based on published general guidelines and are not specific to treatment with BAVENCIO® (avelumab) and INLYTA® (axitinib). You can find a list of possible serious side effects in the Important Safety Information on pages 8-13 of this brochure.

Not all side effects are manageable. If you're having side effects, your doctor may decide to adjust or stop your treatment. That's why it's important to tell your healthcare provider about ANY side effects you are experiencing.



Diarrhea

Diarrhea is defined as having loose or watery stools/bowel movements 3 or more times in 1 day. Call your healthcare provider if you have diarrhea.

Your healthcare provider may recommend that you:

- Try yogurt containing probiotics
- Eat small, frequent meals and foods containing soluble fiber, which can be found in oatmeal, beans, and apples
- Avoid spicy foods, fatty foods, caffeine, and raw fruit
- Drink fluids, such as water, juice, or broth

Diarrhea can be the symptom of a more severe medical condition, so it's important to call your healthcare provider right away. He or she may give you ways to help manage it and may adjust or stop your treatment if needed.



High blood pressure

BAVENCIO with INLYTA may cause your blood pressure to rise. In an INLYTA clinical trial, hypertension occurred as early as 4 days into treatment. On average, this increase was seen within the first month of treatment.

Follow your healthcare provider's instructions about having your blood pressure checked regularly. Call your healthcare provider if your blood pressure is high.

Your healthcare provider may recommend that you:

- Get regular exercise
- Keep your weight under control and lower the amount of sodium you eat in your food
- Lower your alcohol intake

If you develop blood pressure problems, your healthcare provider may lower your dose, stop your treatment, or prescribe medicine to treat your high blood pressure. Tell your healthcare provider if you have high blood pressure or a history of heart disease.

Tips to Help Manage Certain Side Effects (cont'd)



Feeling tired

While on treatment, you may feel tired. Call your healthcare provider for a full evaluation if you feel tired while on your treatment.

If tiredness is a recurring problem, your healthcare provider may recommend that you:

- Take short naps and breaks instead of long ones
- Eat well and drink plenty of fluids
- Stay as active as possible
- Try to maintain normal work and social schedules



Nausea

It is best to call your healthcare provider at the first sign of nausea. Nausea is defined as feeling as if you are about to throw up. It is especially important to speak with your healthcare provider if these symptoms keep you from taking your treatment. Your healthcare provider may prescribe medicine for these symptoms.

Your healthcare provider may recommend that you:

- Eat smaller, more frequent meals
- Avoid fatty, fried, spicy, or highly sweet foods
- Eat bland foods at room temperature and drink clear liquids
- If you vomit, start with small amounts of water, broth, or other clear liquids when you are ready to eat again. If that stays down, then try soft foods. Some examples include gelatin, pudding, yogurt, strained soup, or strained cooked cereal. Slowly work up to eating solid food. Make sure that you do not eat any food that you are allergic to

Let your healthcare provider know right away if you're experiencing any side effect. He or she may give you ways to help manage it and may adjust or stop your treatment if needed.

Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for **BAVENCIO** and full [Prescribing Information/Patient Information](#) for **INLYTA**.



Tips to Help Manage Certain Side Effects (cont'd)



Mouth sores

While on your treatment, you may develop mouth sores. If you develop mouth sores, tell your healthcare provider immediately. He or she may tell you to stop your treatment.

Your healthcare provider may recommend that you:

- Eat foods that are soft, and avoid hot, spicy, or acidic foods
- Use a straw for drinking liquids
- Use an alcohol-free mouthwash, and rinse your mouth often with water mixed with baking soda or salt
- Avoid toothpastes with whiteners (for example, peroxide) and use a soft toothbrush
- Ask your healthcare provider if there are over-the-counter or prescription medicines that may help you manage your condition



Skin conditions

Some patients may have blisters or a rash on the palms of their hands or soles of their feet. This is called hand-foot syndrome. Tell your healthcare provider if you start to develop skin problems. He or she may give you specific treatments, which may include lotions, moisturizers, or pain medicines.

Your healthcare provider may recommend that you:

- Wear loose clothes and avoid tight-fitting shoes and jewelry that rub or chafe the hands and feet
- Use sunscreen
- Clean hands and feet with lukewarm water and gently pat dry
- Apply creams containing urea to the hands and feet liberally and often

Let your healthcare provider know right away if you're experiencing any side effect. He or she may give you ways to help manage it and may adjust or stop your treatment if needed.

Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for **BAVENCIO** and full [Prescribing Information/Patient Information](#) for **INLYTA**.



Tips to Help Manage Certain Side Effects (cont'd)



Hoarseness

Hoarseness is when you have a weak, rough, or harsh voice.

Your healthcare provider may recommend that you:

- Drink plenty of water and avoid irritants like dust, smoke, alcohol, or industrial chemicals
- Write things down to give your voice a break
- Remember to avoid shouting or whispering



Decreased appetite

While on treatment, you may have less desire to eat. But maintaining good nutrition and a healthy weight are important to your overall health. Protein and calories are vital to someone with cancer.

Your healthcare provider may recommend that you:

- Eat when you are hungry and eat several small meals a day—5 or 6 isn't out of the question
- Enjoy snacks and try to make them nutritious. Find calories and protein in dried fruits, nuts, cheeses, and even milkshakes
- Add gravy, butter, or cheese to your favorite foods for added protein and calories
- Drink fluids between meals rather than filling up with them during meals
- If the smell of a certain food is bothering you, it may help to eat it cold or at room temperature
- If you're having trouble tasting your favorite foods, try adding herbs or condiments

Let your healthcare provider know right away if you're experiencing any side effect. He or she may give you ways to help manage it and may adjust or stop your treatment if needed.



Taking BAVENCIO® (avelumab) With INLYTA® (axitinib)

BAVENCIO and INLYTA are each taken in a different way.

BAVENCIO

BAVENCIO is administered by a doctor or a nurse in an infusion center, hospital, or doctor's office.

- Your healthcare provider will give you BAVENCIO via IV infusion over 60 minutes
- Your healthcare provider will give you medicines (an antihistamine and acetaminophen) before the first 4 infusions and subsequently as needed to help reduce infusion reactions
- BAVENCIO is usually given every 2 weeks
- Your healthcare provider will decide how many treatments you need
- Your healthcare provider will do blood tests to check you for certain side effects
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment



On BAVENCIO treatment days, you'll spend time at the infusion center, hospital, or doctor's office. Bring a book, tablet, or your favorite music to pass the time.

INLYTA

Tips to help you when taking INLYTA:

- INLYTA comes in 1-mg and 5-mg tablets taken by mouth. Your healthcare provider will tell you which tablets you should take
- Take INLYTA exactly as prescribed by your healthcare provider
- Your healthcare provider may change your dose if needed
- INLYTA can be taken with or without food
- Take INLYTA 2 times a day approximately 12 hours apart
- Swallow INLYTA tablets whole with a glass of water
- Your healthcare provider should check your blood pressure regularly during treatment with INLYTA
- If you vomit or miss a dose of INLYTA, take your next dose at your regular time. Do not take 2 doses at the same time
- If you take too much INLYTA, call your healthcare provider or go to the nearest hospital emergency room right away
- Do not drink grapefruit juice or eat grapefruit. These may increase the amount of INLYTA in your blood
- Patients taking INLYTA should also avoid St. John's wort. It may reduce the amount of INLYTA in your blood

Starting a treatment plan can take some getting used to. If you have questions, ask your healthcare provider. Keeping note of treatment days may help you remember. Try printing out and using the **Treatment Calendar** on the next page.





Treatment Calendar for BAVENCIO® (avelumab) With INLYTA® (axitinib)

Keeping your treatment schedule organized can help you and your healthcare provider. Use the removable, reusable static cling **Treatment Calendar** at the right with a dry erase pen. Stick it on a flat, smooth surface like a window or a fridge.

Step 1: Write down the day of your first infusion with BAVENCIO. Two weeks from this date, mark the calendar with your next BAVENCIO treatment date.

Step 2: Once you've noted the first day of your BAVENCIO treatment, **mark** each day that you take INLYTA over the next 2 weeks.

Step 3: INLYTA must be taken 2 times a day. **Check** the AM box every morning and the PM box every evening when you take INLYTA.

Example:

	SUN	MON	TUES	WED	THU	FRI	SAT
Date		MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY
BAVENCIO	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>
Date	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY	MM/DD/YY
BAVENCIO	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>	AM <input checked="" type="checkbox"/> PM <input checked="" type="checkbox"/>

Treatment Calendar for BAVENCIO® (avelumab) With INLYTA® (axitinib)

Print out this calendar and follow the steps below.

Step 1: Write down the day of your first infusion with BAVENCIO. Two weeks from this date, **mark** the calendar with your next BAVENCIO treatment date.

Step 2: Once you've noted the first day of your BAVENCIO treatment, **mark** each day that you take INLYTA over the next 2 weeks.

Step 3: INLYTA must be taken 2 times a day. **Check** the AM box every morning and the PM box every evening when you take INLYTA.

	SUN	MON	TUES	WED	THU	FRI	SAT
Date							
BAVENCIO	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>

Date							
BAVENCIO	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>

Date							
BAVENCIO	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>

Date							
BAVENCIO	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
INLYTA	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>	AM <input type="checkbox"/> PM <input type="checkbox"/>

Please see Important Safety Information, full [Prescribing Information](#) and [Medication Guide](#) for BAVENCIO, and full [Prescribing Information and Patient Information](#) for INLYTA at [BavencioPlusInlyta.com/patient](#).



Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for BAVENCIO and full [Prescribing Information/Patient Information](#) for INLYTA.

Questions for Your Healthcare Provider

You and your loved ones will likely have questions about your treatment. Be sure to reach out to your healthcare provider with any concerns or questions you may have. Here are some questions you may find helpful to print out and ask:

Why do I need both medicines?

Is there anything I should do before treatment?

What can I expect during treatment?

How will I know if the treatment is working?

What are common side effects of these 2 medicines? Do you have any tips to help manage them?
Whom do I contact if I have side effects?

I read that I should avoid grapefruit, grapefruit juice, and St. John's wort while taking INLYTA.
Is there anything else I should avoid?

What support resources are available?

Are there any support programs or groups I can join?

Write down any other questions you may have:



Patient Support

As you or your loved one gets started on treatment, you may be looking for some support. Our programs are here to help you get access to treatment. BAVENCIO® (avelumab) with INLYTA® (axitinib) has 2 different support programs, one for each medicine.

Support for eligible patients prescribed BAVENCIO® (avelumab)

CoverOne®

CoverOne® provides eligible patients access and reimbursement support services.

We recognize that each patient’s situation is different, and are dedicated to helping eligible BAVENCIO patients one at a time.

Please contact us at **1-844-8COVER1** if you have any questions.



Phone: 1-844-8COVER1 (1-844-826-8371)
Monday–Friday, 8:00 AM–8:00 PM ET



Web: Visit CoverOne.com

Support for eligible patients prescribed INLYTA® (axitinib)

Pfizer Oncology together™

We’re committed to helping you get your prescribed Pfizer Oncology medicine. Pfizer Oncology Together can help you understand your insurance and identify what financial support may be available for your INLYTA prescription. In addition, we can help find a specialty pharmacy that can fill your prescription.

For live, personalized support



Call **1-877-744-5675** (Monday–Friday, 8:00 AM–8:00 PM ET)

Visit PfizerOncologyTogether.com



Your Oncology Team at a Glance

Print out this page so you can keep a list of your important contacts together in one place for easy access.

Healthcare provider _____

Address _____

Phone _____

Email _____

Other healthcare team member _____

Address _____

Phone _____

Email _____

Notes _____

Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for **BAVENCIO** and full [Prescribing Information/Patient Information](#) for **INLYTA**.

Wallet Card

Download and print out this wallet card at [BavencioPlusInlyta.com/patient](#). Remember to keep it with you at all times.



Notes _____



For more information, ask your healthcare provider
about **BAVENCIO** with **INLYTA** and visit
BavencioPlusInlyta.com/patient



INDICATION

BAVENCIO is a prescription medicine used to treat a type of kidney cancer called renal cell carcinoma (RCC). BAVENCIO may be used with the medicine INLYTA as your first treatment when your kidney cancer has spread or cannot be removed by surgery (advanced RCC).

It is not known if BAVENCIO and INLYTA are safe and effective in children.



Copyright © 2020 EMD Serono, Inc. All rights reserved. March 2020
US/AVE/0619/0228c

Please see Important Safety Information on pages 8-13. Please see full [Prescribing Information](#) and [Medication Guide](#) for **BAVENCIO** and full [Prescribing Information/Patient Information](#) for **INLYTA**.