



For adult patients with advanced urothelial carcinoma (UC), a type of cancer in the bladder or urinary tract, whose cancer has responded or stabilized after platinum-containing chemotherapy

Time for More Moments

There's the possibility for more with BAVENCIO

In a clinical trial for maintenance treatment for advanced urothelial carcinoma, a type of cancer in the bladder or urinary tract, BAVENCIO plus supportive care* showed an increased median overall survival of 21.4 months vs 14.3 months with supportive care alone.

*Supportive care may have included antibiotics, nutritional support, hydration, and pain management.

SELECT SAFETY INFORMATION

BAVENCIO® (avelumab) is a medicine that may treat certain cancers by working with your immune system. BAVENCIO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Please see Select Safety Information throughout and full [Prescribing Information](#) and [Medication Guide](#).



Not an actual patient. BAVENCIO is indicated only for adult patients.

How is advanced bladder cancer treated?

There are 5 common treatments for advanced **bladder cancer**.
You and your doctor can talk about which treatment approach is best for you

Surgery: When doctors do a procedure to take the cancer cells out of your body

Radiation therapy: Strong energy waves are used to kill cancer cells

Chemotherapy: Medicine that stops cancer cells from growing. It can either kill the cancer cells or stop them from dividing. A common type of chemotherapy for advanced bladder cancer is **platinum-containing chemotherapy (PCC)**, such as cisplatin or carboplatin, in combination with gemcitabine

Antibody-drug conjugates: This connects a chemotherapy drug to an antibody, which is a molecule that helps deliver the medicine to the right place in the body

Immunotherapy: A medicine that helps your body's immune system fight and kill cancer cells

Your doctor might also suggest joining a **clinical trial** to try new treatments.

The role of maintenance therapy

- Maintenance therapy may help keep cancer from coming back after treatment
- It may help patients live longer
- It is usually immunotherapy after treatment with chemotherapy



Not sure what a teal-colored word in this brochure means?

Visit [BAVENCIO.com](https://www.bavencio.com) to learn more

SELECT SAFETY INFORMATION (continued)

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with BAVENCIO. Your healthcare provider may treat you with corticosteroids or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with BAVENCIO if you have severe side effects.

Please see **Select Safety Information throughout and full [Prescribing Information and Medication Guide](#).**


 **BAVENCIO**[®]
avelumab
Injection
20 mg/mL

BAVENCIO is an immunotherapy drug— a maintenance treatment that helps your immune system find and help attack cancer cells in advanced bladder cancer

BAVENCIO is used as a maintenance treatment. A maintenance treatment may help keep your cancer from coming back or growing.

BAVENCIO is used after treatment with PCC. PCC is a widely used approach for advanced bladder cancer that has been around for decades and is well understood.

Your immune system can help you fight cancer cells by attacking them with **T cells**, a type of white blood cell. But cancer cells can trick T cells with **PD-L1** by binding PD-1 on a T cell and making it think that the cancer cell is a normal part of the body. **BAVENCIO may help block this connection between PD-L1 and PD-1 so that your body's T cells can fight the cancer cells.**



*This is a general example of how BAVENCIO works. Each patient is different and clinical results may vary. BAVENCIO can cause your immune system to attack normal organs and tissues and can affect the way they work. These problems can sometimes become serious or life threatening and can lead to death.

Please see [Select Safety Information](#) throughout and full [Prescribing Information](#) and [Medication Guide](#).

 **BAVENCIO**[®]
Injection
avelumab 20 mg/mL

Study design

BAVENCIO was studied in 700 patients with advanced bladder cancer that did not progress after **platinum-containing chemotherapy (PCC)**

To be part of the study, patients had to:

- Have locally advanced or metastatic bladder cancer
- Have at least 1 lesion, or damaged area, that was evaluated
- Have gone through 4 to 6 cycles of PCC
- Showed they are physically functional
- Not had any autoimmune diseases (conditions where the immune system attacks your own body) or need a treatment that weakened their immune system

The study measured **overall survival**, or the total time that patients lived after starting in the trial. Patients were given either:

BAVENCIO + supportive care

350 patients

Only supportive care

350 patients

Supportive care for advanced bladder cancer may have included antibiotics to kill bacteria, nutritional support for food and energy, hydration to get enough water, and pain management.

SELECT SAFETY INFORMATION (continued)

BAVENCIO can cause serious side effects. Call or see your healthcare provider right away if you get any new or worsening signs or symptoms, including:

Lung problems

- cough
- shortness of breath
- chest pain

Intestinal problems

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdomen) pain or tenderness

Please see Select Safety Information throughout and full [Prescribing Information and Medication Guide](#).



Not an actual patient.

BAVENCIO[®]
avelumab
Injection
20 mg/mL

Study results

BAVENCIO plus supportive care helped some patients live longer than those on supportive care alone

In the main results of the study:

Patients on **BAVENCIO plus supportive care lived for a median*** of **7.1 months longer** than patients on supportive care alone.

The median overall survival for BAVENCIO plus the best supportive care was 21.4 months and 14.3 months for the best supportive care alone.




BAVENCIO plus supportive care decreased the risk of death by 31% compared with supportive care alone in patients who are stable after receiving PCC.

*The median number is the one that is in the middle of all the results. Half of the people who had BAVENCIO and supportive care lived for more than 21.4 months, and half lived for less.

In the extended study results:

BAVENCIO showed consistent results when the study kept track of patients for over 3 years.



Patients who had **BAVENCIO plus supportive care lived for a median of 8.8 months longer** than those who had only supportive care.

Please see [Select Safety Information](#) throughout and full [Prescribing Information and Medication Guide](#).

Safety

The safety of BAVENCIO was studied in one of the largest trials of patients with advanced bladder cancer—giving you and your doctor the right information to help support educated treatment decisions



Not actual patients.

Almost **9 out of 10 people (88%)** in the study were **able to stay on treatment** and didn't have to permanently stop treatment because of side effects.



The types of side effects seen with BAVENCIO were the same for more than 3 years, and no new side effects were seen during this time.

Please see [Select Safety Information](#) throughout and full [Prescribing Information and Medication Guide](#).

 **BAVENCIO**[®]
avelumab Injection
20 mg/mL

Safety (continued)

Possible side effects

The most common side effects of BAVENCIO include:



Feeling tired



Muscle and bone pain



Urinary tract infection



Rash

Getting medical attention right away may help keep these problems from becoming more serious. Your doctor will check you for these problems during your treatment with BAVENCIO. Your doctor may treat you with corticosteroid or hormone replacement medicines.



Please see [Select Safety Information](#) throughout and full [Prescribing Information](#) and [Medication Guide](#).

BAVENCIO can cause serious side effects.

Call or see your healthcare provider right away if you get any new or worsening signs or symptoms, including:



Lung Problems



Hormone Gland Problems



Intestinal Problems



Kidney Problems



Liver Problems



Skin Problems

Additional serious side effects for BAVENCIO include:



Problems Can Also Happen in Other Organs and Tissues



Infusion-Related Reactions Can Sometimes Be Severe or Life-Threatening



Complications, Including Graft-Versus-Host-Disease (GVHD), in People Who Have Received a Bone Marrow (Stem Cell) Transplant That Uses Donor Stem Cells (Allogeneic)

Tips for your infusion

Remember to note any supplements or over-the-counter medications you may be taking

Also bring fun things to keep you busy, like a book, device, or music.

These steps may help during your infusion:

- ✓ Stay hydrated by drinking plenty of water
- ✓ Wear comfortable, loose-fitting clothing
- ✓ Bring a list of all the medications you currently take
- ✓ Write down any questions you have for your healthcare team
- ✓ Bring something to keep you entertained



SELECT SAFETY INFORMATION (continued)

BAVENCIO can cause serious side effects. Call or see your healthcare provider right away if you get any new or worsening signs or symptoms, including:

Liver problems

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Please see [Select Safety Information throughout and full Prescribing Information and Medication Guide.](#)

How BAVENCIO is given

BAVENCIO is a **first-line maintenance infusion** treatment for advanced bladder cancer given once every 2 weeks



Intravenous
(IV) infusion



About
60 minutes



Every
2 weeks

See infusion-related reactions below

Not an actual patient.
BAVENCIO is indicated
only for adult patients.



You will take BAVENCIO once every 2 weeks. You and your doctor will decide how many treatments you need.



Your doctor or nurse will give you BAVENCIO at an infusion center, hospital, or in your doctor's office. BAVENCIO is given through an IV infusion in your vein that takes about 1 hour to complete.



Your doctor may need to delay or completely stop treatment with BAVENCIO if you have severe side effects.



Infusion related reactions can sometimes be severe or life-threatening. Signs or symptoms of infusion-related reactions may include: chills or shaking, hives, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back pain, or stomach-area (abdomen) pain.

Please see **Select Safety Information** throughout and full **Prescribing Information** and **Medication Guide**.

 **BAVENCIO**[®]
Injection
20 mg/mL

Financial support for BAVENCIO with CoverOne®

At CoverOne®, we're here to help you understand your insurance coverage, access your prescribed BAVENCIO treatment, and support you throughout your treatment journey

CoverOne® may be able to provide access and reimbursement support services for eligible patients.*

Help Understanding Insurance Coverage

- We're here to help you understand how BAVENCIO may be covered by your insurance plan

Copay Assistance/Financial Assistance

- If you have commercial insurance, you may be eligible to pay as little as \$0* out of pocket for BAVENCIO

Patient Assistance Program/Free Drug Program

- Don't have insurance coverage for BAVENCIO? We may be able to provide the medication at no cost if you meet certain eligibility criteria*

Nursing Support

- Our CoverOne® nurses are available to help answer your questions and provide support throughout your BAVENCIO treatment journey

Please see [Select Safety Information](#) throughout and full [Prescribing Information and Medication Guide](#).



CoverOne®

Call us at 1-844-8COVER1 (1-844-826-8371),
Monday–Friday 8 AM to 8 PM EST
Visit us at www.CoverOne.com

***YOUR VOICE IS IMPORTANT.
WORK WITH YOUR DOCTOR TO DEVELOP
A CARE PLAN THAT MEETS YOUR NEEDS***

*Terms and Conditions apply. Some limitations are required by law. Patients covered by federal or state healthcare programs, including Medicare and Medicaid, are not eligible for assistance.

 **BAVENCIO®**
Injection
avelumab 20 mg/mL

IMPORTANT SAFETY INFORMATION (continued)

BAVENCIO can cause serious side effects. Call or see your healthcare provider right away if you get any new or worsening signs or symptoms, including:

Hormone gland problems

- headache that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

Skin problems

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area
- fever or flu-like symptoms
- swollen lymph nodes

Problems can also happen in other organs and tissues. These are not all of the signs or symptoms of immune system problems that can happen with BAVENCIO. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eye sight
- Persistent or severe muscle pain or weakness, muscle cramp
- Low red blood cells, bruising

Infusion-related reactions can sometimes be severe or life-threatening. Signs and symptoms of infusion-related reactions may include:

- chills or shaking
- hives
- flushing
- shortness of breath or wheezing
- dizziness
- feel like passing out
- fever
- back pain
- stomach-area (abdomen) pain

Complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with BAVENCIO. Your healthcare provider will monitor you for these complications.

Please see full [Prescribing Information](#) and [Medication Guide](#).



IMPORTANT SAFETY INFORMATION (continued)

Before you receive BAVENCIO, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- have heart problems or high blood pressure
- have diabetes
- have a high cholesterol level in your blood
- are pregnant or plan to become pregnant. BAVENCIO can harm your unborn baby

Females who are able to become pregnant:

- You should use an effective method of birth control during your treatment and for at least 1 month after the last dose of BAVENCIO. Talk to your healthcare provider about birth control methods that you can use during this time
- are breastfeeding or plan to breastfeed. It is not known if BAVENCIO passes into your breast milk. Do not breastfeed during treatment and for at least 1 month after the last dose of BAVENCIO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of BAVENCIO as maintenance treatment in people with urothelial carcinoma (UC) whose cancer responded or stabilized after platinum-containing chemotherapy as first treatment include:

- feeling tired
- muscle and bone pain
- urinary tract infection
- rash

The most common side effects of BAVENCIO in people with UC after platinum-containing chemotherapy that did not work, or is no longer working, include:

- feeling tired
- infusion-related reactions including chills, fever, back pain, redness, and shortness of breath
- muscle and bone pain
- nausea
- decreased appetite
- urinary tract infection

These are not all the possible side effects of BAVENCIO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full full [Prescribing Information](#) and [Medication Guide](#).



Talk to your doctor about
BAVENCIO
to learn if there is a

possibility for **MORE**

BAVENCIO first-line maintenance may help make a difference in your treatment journey for advanced bladder cancer



Not actual patients.

Still have questions about BAVENCIO? Talk to your doctor or visit [BAVENCIO.com](https://www.bavencio.com) to learn more

Please see **Select Safety Information** throughout and full **Prescribing Information and Medication Guide**.



© 2025 Merck KGaA, Darmstadt, Germany and/or its affiliates. All rights reserved.
EMD Serono is the Healthcare business of Merck KGaA, Darmstadt, Germany in the U.S. and Canada.
BAVENCIO and CoverOne are registered trademarks of Merck KGaA, Darmstadt, Germany, or its affiliates.
US-AVE-01899 09/25

